

AD-A157 484

THE IONIZATION EQUILIBRIUM OF OPTICALLY THICK ARGON
Z-PINCH PLASMAS FOR E. (U) NAVAL RESEARCH LAB
WASHINGTON DC J P APRUZESE ET AL. 02 JUL 85

1/1

UNCLASSIFIED

NRL-MR-5689


F/G 20/9

NL

END

FILED

DTIC



MICROCOPY RESOLUTION TEST CHART
NBS-1963-A

AD-A157 484

2
NRL Memorandum Report 5609

The Ionization Equilibrium of Optically Thick Argon Z-Pinch Plasmas for Electron Temperatures Between 25 and 65 eV

J. P. APRUZESE, P. C. KEPPLER, AND J. DAVIS

*Plasma Radiation Branch
Plasma Physics Division*

July 2, 1985

This work was supported in part by the Strategic Defense Initiative Organization and the Defense Nuclear Agency under Subtask T99QMXLA, work unit 00004 and work unit title "Advanced Simulation Concepts."



DTIC
ELECTE
JUL 17 1985
S D G

NAVAL RESEARCH LABORATORY
Washington, D.C.

Approved for public release; distribution unlimited.

DTIC FILE COPY

REPORT DOCUMENTATION PAGE				
1a REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b RESTRICTIVE MARKINGS		
2a SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
2b DECLASSIFICATION / DOWNGRADING SCHEDULE				
4 PERFORMING ORGANIZATION REPORT NUMBER(S) NRL Memorandum Report 5609		5 MONITORING ORGANIZATION REPORT NUMBER(S)		
6a NAME OF PERFORMING ORGANIZATION Naval Research Laboratory	6b OFFICE SYMBOL (If applicable) Code 4720	7a NAME OF MONITORING ORGANIZATION Defense Nuclear Agency		
6c ADDRESS (City, State, and ZIP Code) Washington, DC 20375-5000		7b ADDRESS (City, State, and ZIP Code) Washington, DC 20305		
8a NAME OF FUNDING / SPONSORING ORGANIZATION Defense Nuclear Agency	8b OFFICE SYMBOL (If applicable) RAEV	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c ADDRESS (City, State, and ZIP Code) Washington, DC 20305		10 SOURCE OF FUNDING NUMBERS		
		PROGRAM ELEMENT NO. 62715H	PROJECT NO. 85-579	TASK NO. DN880-191
11 TITLE (Include Security Classification) The Ionization Equilibrium of Optically Thick Argon Z-Pinch Plasmas for Electron Temperatures Between 25 and 65 eV				
12 PERSONAL AUTHOR(S) Apruzese, J.P., Kepple, P.C., and Davis, J.				
13a TYPE OF REPORT Interim	13b TIME COVERED FROM TO	14. DATE OF REPORT (Year, Month, Day) 1985 July 2	15. PAGE COUNT 28	
16 SUPPLEMENTARY NOTATION (See page ii)				
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP		
			Ionization equilibrium Argon plasma	
			Gamble-II generator Collisional pumping, <i>gamma</i>	
19 ABSTRACT (Continue on reverse if necessary and identify by block number)				
<p>Calculations of the ionization balance of optically thick argon plasmas of electron temperature 25-65 eV are presented. A multistage, multilevel model of argon ions with accurate radiation transport of ~ 50 lines in the sodium, neon, and fluorine-like stages forms the basis of the calculations. By comparison with similar optically thin calculations, the effects on ionization of radiative and collisional pumping of excited states of these stages is delineated. In particular, the broad temperature distribution of the neon-like stage is compressed by line photon pumping to a narrower regime of existence because of ionization from the excited states. The relevance to the 3p-3s lasing scheme in neon-like argon is briefly discussed. <i>gamma</i></p>				
20 DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a NAME OF RESPONSIBLE INDIVIDUAL Jack Davis			22b TELEPHONE (Include Area Code) (202) 767-3278	22c OFFICE SYMBOL Code 4720

16. SUPPLEMENTARY NOTATION

This work was supported in part by the Strategic Defense Initiative Organization and the Defense Nuclear Agency under Subtask T99QMXLA, work unit 00004 and work unit title "Advanced Simulation Concepts."

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist.	Avail and/or Special
A/1	



CONTENTS

I. INTRODUCTION	1
II. DESCRIPTION OF THE ATOMIC MODEL AND RADIATION TRANSPORT	2
III. RESULTS	4
ACKNOWLEDGMENT	6
REFERENCES	15

THE IONIZATION EQUILIBRIUM OF OPTICALLY THICK ARGON Z-PINCH PLASMAS FOR ELECTRON TEMPERATURES BETWEEN 25 AND 65 eV

I. Introduction

Due to some recent successful experiments at Livermore,^{1,2} which were preceded by a host of theoretical calculations,³⁻¹¹ considerable interest now centers upon neon-like ions in plasmas as very promising for achievement of population inversions leading to lasing in the extreme ultraviolet to x-ray regions of the spectrum. Although the first successful demonstrations used frequency-doubled neodymium laser light at 0.53 μm as a plasma driver, use of Z-pinch plasmas coupled to large pulsed power generators as a lasing medium is also highly attractive due to the large gain lengths (up to 4 cm) and immense energies (~ 1 MJ) available to couple to the plasma. The Gamble-II device at NRL, while of relatively modest energy, has proven to be a very efficient producer of K-shell radiation from neon plasmas^{12,13}. The use of fast plasma opening switches¹⁴ to reduce pulse risetimes has produced more orderly pinches with instabilities considerably less pronounced than those previously observed¹⁵. These results suggest that Gamble-II would be an excellent device to test lasing concepts on a Z-pinch. Argon, stripped to the neon-like stage, is a prime candidate for lasing given both the available machine energy and expectation of high quality pinches. It is the purpose of this Memorandum Report to investigate the plasma conditions (temperature, density, size) which will optimize the production of the neon-like stage and also the pumping of the upper lasing 3p manifold of J-sublevels. We also identify and clarify the physical mechanisms which are responsible for the functional behavior of the ionization balance and pumping of the 3p levels.

II. Description of the Atomic Model and Radiation Transport

The atomic model consists of all ground states of argon with the following excited states in the sodium-like, neon-like, and fluorine-like ions:

	State	Statistical Weight	Energy Above Ground (eV)
Ar VIII (sodium-like)	$3p^2P$	6	17.695
	$3d^2D$	10	41.219
	$4s^2S$	2	71.377
	$4p^2P$	6	77.986
	$4d^2D$	10	86.443
	$4f^2F$	14	88.841
	$[n=5]$	50	103.20
Ar IX (neon-like)	$[2p^5 3s]$	12	252.16
	$[2p^5 3p]$	36	270.18
	$[2p^5 3d]$	60	293.76
	$[2s2p^6 3s]$	4	329.43
	$[2s2p^6 3p]$	12	346.82
	$[2p^5 n=4]$	192	349.81
	$[2s2p^6 3d]$	20	370.19
Ar X (fluorine-like)	$2s2p^6 2s$	2	75.734
	$[2p^4 3s]$	30	283.86
	$[2p^4 3p]$	90	301.96
	$[2p^4 3d]$	150	324.69
	$[2s2p^5 3s]$	24	353.97
	$[2s2p^5 3p]$	72	371.25
	$[2s2p^5 3d]$	120	393.44
	$[2p^4 n=4]$	480	394.20
	$[2s2p^5 n=4]$	384	435.00

The states enclosed in square brackets [] are composites of several states with similar energy.

The rate table associated with this model includes:

1. Photoionization rates from each ground state and each excited state. These rates were calculated using the hydrogenic approximation with Karzas and Latter¹⁶ Gaunt factors, corrected for equivalent electrons.
2. Collisional ionization from each ground and excited state to the ground state above. These rates were calculated using the exchange classical impact parameter technique.¹⁷
3. Dielectronic recombination rates from all ground states to the adjacent ground state. These rates include (implicitly) the effect of the excited states¹⁸.
4. The Einstein A coefficients were calculated by Cowan's¹⁹ code with appropriate averages for composite levels.
5. Collisional excitation rates for dipole allowed transitions were calculated in the semiclassical impact parameter approximation²⁰. For transitions with no dipole allowed component the collisional excitation rates were calculated with a distorted wave²¹ code.
6. Detail balance was used to obtain the radiative recombination rates from the photoionization rates, the 3-body recombination rates from the collisional ionization rates, and the collisional deexcitation rates from the collisional excitation rates.

Radiation transport is calculated by a cell-to-cell photon coupling technique using a matrix C_{ij} for each transition which is the probability that a line photon emitted in cell i is absorbed in cell j . Details of the technique have been given previously.^{22,23} The steady-state solution for the state and level populations, consistent with the radiation field, is obtained by an iteration technique.²⁴

III. Results

Two argon ion densities ($3 \times 10^{18} \text{ cm}^{-3}$ and $6 \times 10^{19} \text{ cm}^{-3}$) have been chosen for the calculations. These densities represent the approximate lower and upper limits of those achieved in imploding neon gas puff experiments on the NRL Gamble-II generator. The 25-65 eV temperature regime is chosen to examine the behavior of the neon-like argon stage. The results of our calculations are presented in Figs. 1-7. In each case the ground state ionic fraction for the neon-, fluorine-, and sodium-like stages is shown as a function of electron temperature. The total ion density and assumed plasma size differ from figure to figure. In Figs. 1-4, an ion density of $6 \times 10^{19} \text{ cm}^{-3}$ is assumed, and in the remainder of the figures the ion density used was $3 \times 10^{18} \text{ cm}^{-3}$. For each chosen ion density, cylindrical plasma diameters of 1.8 mm and 0.5 mm were used, and a calculation was also performed in which all lines were assumed optically thin (equivalent to the zero-size limit). The objectives were to approximate the experimental conditions, expected with argon gas-puff implosions on Gamble II, and, through use of the optically thin calculation, to gauge the effects of radiative pumping on ionization.

Figs. 1 and 2 present the ionization curves for ion density $6 \times 10^{19} \text{ cm}^{-3}$ and cylindrical diameter 1.8 mm. The only difference is that, for Fig. 1, no sodium-like excited states were included in the model. Only a minor effect on the neon-like stage distribution occurs, but note that the sodium-like stage persists at higher temperatures when the calculation excludes excited states for the sodium-like stage. This is because a significant amount of ionization occurs from the excited states, and excluding them from the model inevitably results in a less ionized plasma. When the excited states are included, the fluorine-like stage becomes substantially more prevalent, reflecting the extra ionization. These effects have also been previously noted for K-shell plasmas.²⁵

Figures 2, 3, and 4 display the effects of line opacity on argon plasma ionization, for a fixed total argon ion density of $6 \times 10^{19} \text{ cm}^{-3}$. As the assumed plasma size decreases from 1.8 mm to 0.5 mm (Fig. 2 compared to Fig. 3), little effect is noted. This is due to the fact that the optical depths of the principal resonance lines are already so high (10^2 - 10^3) that

the radiative pumping rate is saturated by collisional destruction of the line photons after a few scatterings in the lines. Therefore, since the line pumping does not change, the ionization does not change. Comparison of Figs. 5 and 6 - where the ion density is $3 \times 10^{18} \text{ cm}^{-3}$ - reveals this same lack of change at lower ion density. However, when the plasma is assumed optically thin in all the lines, substantial reduction of the ionization is observed. This is seen specifically by comparing Fig. 4 to Fig. 3 and Fig. 7 to Fig. 6. When there is no radiative pumping of the excited levels, there is less ionization from those levels and, therefore, a less highly ionized plasma. Experimental achievement of an optically thin argon plasma would require a characteristic size of a few microns - currently unrealizable with a Z-pinch. The optically thin plasmas also exhibit a broader distribution of the neon-like stage. This is a manifestation of the relative lack of excited state populations in these cases, allowing the characteristic "closed shell" properties of the neon-like ground state to produce the expected broad distribution closer to that seen in the coronal limit.²⁶

Further insight into this last point is obtained by comparing Fig. 4 and Fig. 7. Both curves reflect optically thin calculations, the only difference being the ion density. At the lower ion density of Fig. 7, an extremely broad peak in the neon-like distribution is seen, similar to that obtained in the coronal limit by Shull and Van Steenberg²⁶ (also plotted) but at lower temperatures than their results. Figure 4 reveals that at higher densities the peak is narrowed even though the calculation still assumes zero optical depth at all frequencies. The reason for this difference is that excited states are populated collisionally as well as radiatively. Even though no radiative excitations occur in those calculations, the increased collisional population of the excited states at higher ion density removes some of the "closed shell" property from the neon-like stage, resulting in more ionization (from the excited states) and therefore a narrower peak further removed from the coronal limit than at the lower ion density of $3 \times 10^{18} \text{ cm}^{-3}$.

Finally, we address the implications of these results for the achievement of 3p-3s lasing in neon-like Ar IX. At first it would seem that optimum plasma conditions would lie where the neon-like stage

maximizes, i.e. at $T_e = 35-45$ eV. Actually, the optimum temperature for 3p-3s lasing is in the 50-55 eV range. Figure 8 illustrates the reason for this difference. In this figure is plotted the fraction of all particles whose energies lie above a certain threshold for a Maxwellian distribution. For Ar IX, the 3p levels are approximately 270 eV above the ground state. At $T_e = 40$ eV, only $\sim 0.3\%$ of the electrons are capable of exciting these levels. However, at 55 eV $\sim 2\%$ of the electrons have sufficient energy to excite the upper lasing 3p levels. This more than compensates for the reduction in neon-like ground state population. Our calculations indicate that the fluorine-like resonance line complex, $2s^2 2p^5 - 2s^2 2p^4 3d$ at 38.5Å will emit with about the same intensity as the neon-like resonance line complex $2p^6 - 2p^5 3d$ at 42Å when optimal pumping conditions are achieved for the neon-like species.

The present model does not contain enough detail in the neon-like excited state manifold to extract detailed gain behavior, and such calculations are reserved for a future report. The ionization balance calculations, however, are affected little by breaking composite levels into J-sublevels for more detailed gain calculations.

ACKNOWLEDGMENT

This work was supported by the U.S. Strategic Defense Initiative Organization and the Defense Nuclear Agency.




Fig. 1. The ground state ionic fractions for sodium-like, neon-like, and fluorine-like argon are plotted against electron temperature for a total argon ion density of $6 \times 10^{19} \text{ cm}^{-3}$ and cylindrical plasma diameter of 1.8 mm. No sodium-like excited states are included in this calculation.




Fig. 2. As in Fig. 1, except that the sodium-like excited states described in Section II are included in the calculation.




Fig. 3. As in Fig. 2, except that the plasma diameter is assumed to be 0.5 mm.




Fig. 4. As in Figs. 2 and 3, except that the plasma is assumed optically thin (zero size limit).




Fig. 5. The ground state ionic fractions for sodium-like, neon-like, and fluorine-like argon are plotted against electron temperature for a total argon ion density of $3 \times 10^{18} \text{ cm}^{-3}$ and a cylindrical plasma diameter of 1.8 mm.




Fig. 6. As in Fig. 5, except that the plasma diameter is assumed to be 0.5 mm.




Fig. 7. As in Figs. 5 and 6 except that the plasma is assumed to be optically thin (zero size limit). The coronal equilibrium results of Ref. 26 for neon-like argon are also shown.




Fig. 8. The fraction of all particles in a Maxwellian distribution whose energies E lie above a threshold energy E_0 is plotted against ratio of temperature to E_0 .

References

1. M. D. Rosen, et al., Phys. Rev. Lett. 54, 106 (1985).
2. D. L. Matthews, et al., Phys. Rev. Lett. 54, 110 (1985).
3. A. N. Zherikhin, K. N. Koshelev, and V. S. Letokhov, Sov. J. Quantum Electron, 6, 82 (1976).
4. A. V. Vinogradov, I. I. Sobel'man, and E. A. Yukov, Sov. J. Quantum Electron, 7, 32 (1977).
5. L. A. Vainshtein, A. V. Vinogradov, V. I. Safronova, and I. Yu. Skobelev, Sov. J. Quantum Electron, 8, 239 (1978).
6. A. V. Vinogradov and V. N. Shylaptsev, Sov. J. Quantum Electron 13, 1511 (1983).
7. P. L. Hagelstein, Lawrence Livermore National Laboratory Report No. UCRL-53100, 1981.
8. U. Feldman, A. K. Bhatia, and S. Suckewer, J. Appl. Phys. 54, 2188 (1983).
9. J. P. Apruzese and J. Davis, Phys. Rev. A 28, 3686 (1983).
10. U. Feldman, J. F. Seely, and A. K. Bhatia, J. Appl. Phys. 56, 2475 (1984).
11. R. H. Dixon and R. C. Elton, J. Opt. Soc. Am. B 1, 232 (1984).
12. S. J. Stephanakis, et al., IEEE Pub. No. 84CH1958-8, 1984.
13. S. J. Stephanakis, private communication.
14. R. A. Meger, R. J. Comisso, G. Cooperstein, and S. A. Goldstein, Appl. Phys. Lett. 42, 943 (1983).
15. S. J. Stephanakis, S. W. McDonald, R. A. Meger, P. F. Ottinger, F. C. Young, C. G. Mehlman, J. P. Apruzese, and R. E. Terry, Bull. Am. Phys. Soc. 29, 1232 (1984).
16. W. J. Karzas and R. Latter, Astrophys. J. Suppl. Ser. 6, 167 (1961).
17. A. Burgess, in Proceedings of the Symposium on Atomic Collision Processes in Plasmas, Culham, England, Report No. 4818, 63, 1964 (unpublished); and A. Burgess, H. P. Summers, D. M. Cochrane, and R. W. P. McWhirter, Mon. Not. R. Astron. Soc. 179, 275 (1977).
18. V. L. Jacobs, J. Davis, P. C. Kepple and M. Blaha, Astrophys. J. 211, 605 (1977).
19. R. D. Cowan, Phys. Rev. 163, 54 (1967); J. Opt. Soc. Am. 58, 808 (1968).

20. J. Davis, J. Quant. Spectros. Radiat. Transfer, 14, 549 (1974).
21. J. Davis, P. C. Kepple, and M. Blaha, J. Quant. Spectrosc. Radiat. Transfer 16, 1043 (1976).
22. J. P. Apruzese, J. Davis, D. Duston, and K. G. Whitney, J. Quant. Spectrosc. Radiat. Transfer 23, 479 (1980).
23. J. P. Apruzese, J. Quant. Spectrosc. Radiat. Transfer 25, 419 (1981).
24. J. P. Apruzese, J. Davis, D. Duston, and R. W. Clark, Phys. Rev. A 29, 246 (1984).
25. J. P. Apruzese and J. Davis, Phys. Rev. A 31, 2976 (1985).
26. J. M. Shull and M. Van Steenberg, Astrophys. J. Suppl. Ser. 48, 95 (1982).

DISTRIBUTION LIST

Assistant to the Secretary of Defense Atomic Energy Washington, D.C. 20301 ATTN: Executive Assistant	1 Copy
Defense Technical Information Center Cameron Station 5010 Duke Street Alexandria, Va 22314	2 copies
Director Defense Intelligence Agency Washington, D.C. 20301 ATTN: DT-1B R. Rubenstein	1 Copy
Director Defense Nuclear Agency Washington, D.C. 20305 ATTN: DDST ATTN: TITL ATTN: RAEV ATTN: STVI	1 copy 4 copies 1 copy 1 copy
Commander Field Command Defense Nuclear Agency Kirtland AFB, New Mexico 87115 ATTN: FCPR	1 Copy
Chief Field Command Livermore Division Department of Defense P.O. Box 808 Livermore, CA 94550 ATTN: FCPRL	1 Copy
Director Joint Strat TGT Planning Staff Offutt AFB Omaha, Nebraska 68113 ATTN: JSAS	1 Copy
Undersecretary of Defense for RSCH and ENGRG Department of Defense Washington, D.C. 20301 ATTN: Strategic and Space Systems (OS)	1 Copy

Deputy Chief of Staff for RSCH DEV and ACQ
Department of the Army
Washington, D.C. 20301
ATTN: DAMA-CSS-N

1 Copy

Commander
Harry Diamond Laboratories
Department of the Army
2800 Powder Mill Road
Adelphi, MD 20783
ATTN: DELHD-N-NP
ATTN: DELHD-R J. Rosado
ATTN: DELHD-TA-L (Tech. Lib.)

1 copy each

U.S. Army Missile Command
Redstone Scientific Information Center
Attn: DRSMI-RPRD (Documents)
Redstone Arsenal, Alabama 35809

3 Copies

Commander
U.S. Army Missile Command
Redstone Arsenal, Alabama 35898
ATTN: DRCPM-PE-EA

1 copy

Commander
U.S. Army Nuclear and Chemical Agency
7500 Backlick Road
Building 2073
Springfield, VA 22150
ATTN: Library

1 copy

Commander
Naval Intelligence Support Center
4301 Suitland Road, Bldg. 5
Washington, D.C. 20390
ATTN: NISC-45

1 Copy

Commander
Naval Weapons Center
China Lake, California 93555
ATTN: Code 233 (Tech. Lib.)

1 Copy

Officer in Charge
White Oak Laboratory
Naval Surface Weapons Center
Silver Spring, Md. 20910
ATTN: Code R40
ATTN: Code F31

1 Copy each

Air Force Weapons Laboratory Kirtland AFB, New Mexico 87117 ATTN: SUL ATTN: CA ATTN: APL ATTN: Lt. Col Generosa	1 Copy each
Deputy Chief of Staff Research, Development and Accounting Department of the Air Force Washington, D. C. 20330 ATTN: AFRDQSM	1 Copy
Commander U.S. Army Test and Evaluation Command Aberdeen Proving Ground, MD 21005 ATTN: DRSTE-EL	1 Copy
Space and Missile Systems Organization/SK Air Force Systems Command Post Office Box 92960 Worldway Postal Center Los Angeles, CA 90009 ATTN: SKF P. Stadler (Space Comm. Systems)	1 Copy
AVCO Research and Systems Group 201 Lowell Street Wilmington, MA 01887 ATTN: Library A830	1 Copy
BDM Corporation 7915 Jones Branch Drive McLean, Virginia 22101 ATTN: Corporate Library	1 Copy
Berkeley Research Associates P.O. Box 983 Berkeley, CA 94701 ATTN: Dr. Joseph Workman	1 Copy
Berkeley Research Associates P.O. Box 852 5532 Hempstead Way Springfield, VA 22151 ATTN: Dr. Joseph Orens ATTN: Dr. Nino Pereira	1 Copy each
Boeing Company P. O. Box 3707 Seattle, WA 98134 ATTN: Aerospace Library	1 Copy

The Dikewood Corporation 1613 University Bldv., N.E. Albuquerque, New Mexico 8710 ATTN: L. Wayne Davis	1 Copy
EG and G Washington Analytical Services Center, Inc. P. O. Box 10218 Albuquerque, New Mexico 87114 ATTN: Library	1 Copy
General Electric Company Space Division Valley Forge Space Center P. O. Box 8555 Philadelphia, PA 19101 ATTN: J. Peden	1 Copy
General Electric Company - Tempo Center for Advanced Studies 816 State Street P.O. Drawer QQ Santa Barbara, CA 93102 ATTN: DASIAC	1 Copy
Institute for Defense Analyses 1801 N. Beauregard St. Alexandria, VA 22311 ATTN: Classified Library	1 Copy
IRT Corporation P.O. Box 81087 San Diego, CA 92138 ATTN: R. Mertz	1 Copy
JAYCOR 11011 Forreyane Rd. P.O. Box 85154 San Diego, CA 92138 ATTN: E. Wenaas	1 Copy
JAYCOR 205 S. Whiting Street, Suite 500 Alexandria, VA 22304 ATTN: R. Sullivan	1 Copy
KAMAN Sciences Corp. P. O. Box 7463 Colorado Springs, CO 80933 ATTN: J. Hoffman ATTN: A. Bridges ATTN: D. Bryce ATTN: W. Ware	1 copy each

Lawrence Livermore National Laboratory University of California P.O. Box 808 Livermore, California 94550 Attn: DOC CDN for L-153 Attn: DOC CDN for L-47 L. Wouters Attn: DOC CDN for Tech. Infor. Dept. Lib.	1 copy each
Lockheed Missiles and Space Co., Inc. P. O. Box 504 Sunnyvale, CA 94086 Attn: S. Taimlty Attn: J.D. Weisner	1 copy each
Lockheed Missiles and Space Co., Inc. 3251 Hanover Street Palo Alto, CA 94304 Attn: J. Perez	1 Copy
Maxwell Laboratory, Inc. 9244 Balboa Avenue San Diego, CA 92123 ATTN: A. Kolb ATTN: M. Montgomery ATTN: J. Shannon	1 Copy each
McDonnell Douglas Corp. 5301 Bolsa Avenue Huntington Beach, CA 92647 ATTN: S. Schneider	1 Copy
Mission Research Corp. P. O. Drawer 719 Santa Barbara, CA 93102 ATTN: C. Longmire ATTN: W. Hart	1 Copy each
Mission Research Corp.-San Diego 5434 Ruffin Rd. San Diego, California 92123 ATTN: Victor J. Van Lint	1 Copy
Northrop Corporation Northrop Research and Technology Center 1 Research Park Palos Verdes Peninsula, CA 90274 ATTN: Library	1 Copy
Northrop Corporation Electronic Division 2301 120th Street Hawthorne, CA 90250 ATTN: V. Damarting	1 Copy

Physics International Company 2700 Merced Street San Leandro, CA 94577 Attn: C. Stallings Attn: C. Gilman	1 Copy each
R and D Associates P.O. Box 9695 Marina Del Rey, CA 90291 ATTN: W. Graham, Jr. ATTN: P. Haas	1 Copy each
Sandia National Laboratories P.O. Box 5800 Albuquerque, New Mexico 87115 ATTN: Doc Con For 3141 ATTN: D. McDaniel ATTN: P. VanDevender ATTN: K. Matzen, Code 4247	1 copy each
Science Applications, Inc. P. O. Box 2351 La Jolla, CA 92038 ATTN: R. Beyster	1 copy
Spire Corporation P. O. Box D Bedford, MA 01730 ATTN: R. Little	1 copy
SRI International 333 Ravenswood Avenue Menlo Park, CA 94025 ATTN: S. Dairiki	1 copy
S-CUBED P. O. Box 1620 La Jolla, CA 92038 ATTN: A. Wilson	1 copy
Director Strategic Defense Initiative Organization Pentagon 20301-7100 ATTN: Lt. Col Richard Gullickson/DEO Dr. Dwight Duston	1 copy each
Texas Tech University P.O. Box 5404 North College Station Lubbock, TX 79417 ATTN: T. Simpson	1 copy

TRW Defense and Space Systems Group
One Space Park
Redondo Beach, CA 90278
ATTN: Technical Information Center

1 Copy

Vought Corporation
Michigan Division
38111 Van Dyke Road
Sterling Heights, Maine 48077
ATTN: Technical Information Center
(Formerly LTV Aerospace Corp.)

1 Copy

Naval Research Laboratory
Plasma Radiation Branch
Washington, D.C. 20375

Code 4720	-	50	Copies
Code 4700	-	26	Copies
Code 2628	-	20	Copies

Director of Research
U.S. Naval Academy
Annapolis, MD 21402

2 Copies

DEPARTMENT OF THE NAVY

NAVAL RESEARCH LABORATORY
Washington, D.C. 20375-5000

OFFICIAL BUSINESS

PENALTY FOR PRIVATE USE, \$300



POSTAGE AND FEES PAID
DEPARTMENT OF THE NAVY
DoD-316
THIRD CLASS MAIL



END

FILMED

9-85

DTIC